

A Canonical Relativistic Approach to Quantize Electromagnetic Field in the Presence of Moving Magneto-Dielectric Media

M. Amooshahi¹ ^{*}

¹ Faculty of science, University of Isfahan ,Hezar Jarib Ave., Isfahan,Iran

August 9, 2009

Abstract

A canonical relativistic formulation is introduced to quantize electromagnetic field in the presence of a polarizable and magnetizable moving medium. The medium is modeled by a continuum of the second rank antisymmetric tensors in a phenomenological way. The covariant wave equation for the vector potential and the covariant constitutive equation of the medium are obtained as the Euler- Lagrange equations using the Lagrangian of the total system. A fourth rank tensor which couples the electromagnetic field and the medium is introduced. The susceptibility tensor of the medium is obtained in terms of this coupling tensor. The noise polarization tensor is calculated in terms of both the coupling tensor and the ladder operators of the tensors modeling the medium.

PACS No: 12.20.Ds, 42.50.Nn

Key words: Canonical field quantization, Magnetodielectric medium, Susceptibility tensor, Coupling tensor, Covariant constitutive equation

^{*}Amooshahi@sci.ui.ac.ir

[†]Tel: 98-311-793-2418, Fax: 98-311-793-2409

1 Introduction

The electromagnetic field can be quantized in the presence of an absorbing polarizable and magnetizable medium by modeling such a medium by two independent collections of oscillators[1, 2]. This method, in addition to its simplicity and beauty, covers some general cases. This quantization scheme has been generalized for anisotropic media and for spatially and temporarily dispersive media [3, 4]. It has been shown that this formalism can be obtained canonically using a Lagrangian function[5]. The quantization method is also applicable for a non-linear magnetodielectric medium. One advantage of this method is that, in contrast of the Huttner-Barnett model [6, 7], the electric and magnetic polarization fields are not needed to be included in the Lagrangian of the total system as a part of the degrees of freedom of the medium. That is, the oscillators modeling the medium constitute solely the degrees of freedom of the medium and are able to describe both polarizability and absorptivity of the medium.

The electromagnetic field quantization in the presence of an anisotropic dielectric media with spatio-temporal dispersion has also been treated previously in the base of the quantum damped polariton model [8].

In the present work the previous model [1]-[5] has been generalized for a moving medium by using a relativistically covariant approach. The medium is modeled, in a phenomenological way, by a continuum of the second rank antisymmetric tensors labeled by a continuous parameter ω . The tensors modeling the medium describe both polarizability and absorptivity of the medium. In order to represent a canonical quantization in a relativistically covariant language, we apply the Lorentz gauge and the Gupta-Bleuler method is used [9]. In Gupta-Bleuler quantization formalism the components of the four-vector potential A^μ constitute the dynamical variables of electromagnetic field and only those vectors, $|\psi\rangle$ in the Hilbert space of the total system, are admitted for which the expectation value of the gauge condition, $\langle\psi|\partial_\mu A^\mu|\psi\rangle = 0$, is satisfied. In this formalism the Lorentz condition can not be carried over to the field operators A^μ as $\partial_\mu A^\mu = 0$, since the condition $\partial_\mu A^\mu = 0$ as an operator identity is incompatible to the commutation relations of the operator A^μ and its conjugate dynamical variable.

For simplicity we set $c = 1$ for the speed of light in the vacuum and choose the space-time metric as $g^{\mu\nu} = \text{dia}(-1, -1, -1, 1)$.

2 Relativistic classical electrodynamics in the presence a magnetodielectric medium

In order to have a classical treatment of electrodynamics in the presence of a moving polarizable and magnetizable medium , the medium is modeled by a continuum of the second rank antisymmetric tensors $Y^{\alpha\beta}(\omega, x)$ labeled by a continuous parameter ω . The Lagrangian density of the total system is proposed as

$$\begin{aligned} \mathcal{L}(x) = & \frac{1}{2} \partial_\mu A_\nu \partial^\mu A^\nu - \int_0^\infty d\omega f^{\mu\nu\alpha\beta}(\omega, x) \partial_\mu A_\nu Y_{\alpha\beta}(\omega, x) \\ & + \frac{1}{2} \int_0^\infty d\omega \left[\partial_\mu Y_{\alpha\beta}(\omega, x) \partial^\mu Y^{\alpha\beta}(\omega, x) - \omega^2 Y_{\alpha\beta}(\omega, x) Y^{\alpha\beta}(\omega, x) \right] \end{aligned} \quad (1)$$

where $x = (t, \vec{x})$ stands for the coordinates of the space-time and A^μ is the four vector potential . Here the tensor $f^{\mu\nu\alpha\beta}(\omega, x)$ is called the coupling tensor of electromagnetic field and the medium and is antisymmetric with respect to the first two indices μ, ν and also with respect to the last two indices α, β . Using the Lagrangian (1), the Euler- Lagrange equations for the vector potential are obtained as

$$\square^2 A^\nu = 4\pi \partial_\mu K^{\mu\nu} \quad (2)$$

where $\square^2 = \frac{\partial^2}{\partial t^2} - \nabla^2$ is the d'Alembert's operator and

$$K^{\mu\nu}(x) = \frac{1}{4\pi} \int_0^\infty d\omega f^{\mu\nu\alpha\beta}(\omega, x) Y_{\alpha\beta}(\omega, x) \quad (3)$$

is the antisymmetric electric-magnetic polarization tensor of the medium. Accordingly, the polarization of the medium can be defined naturally in terms of the antisymmetric tensors $Y^{\alpha\beta}(\omega, x)$ modeling the medium and the polarization fields of the medium are not needed to be included explicitly in the Lagrangian of the total system. The electric-magnetic polarization tensor $K^{\mu\nu}$ is related to the field strength tensor $F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu$ by the covariant form of the constitutive equation of the moving medium. Before obtaining this equation from the Euler- Lagrange equations, it is necessary to point out the following remarks:

1. The electric polarization density of a moving medium depends on both electric and magnetic field , even though the electric polarization is dependent only on the electric field when the medium is at rest. This is so, since both

the magnetic and electric forces affect the electric polarization of the medium when it moves. Similarly the magnetic polarization of a moving medium in general may be dependent on both electric and magnetic field[10, 11].

2. A moving medium may be anisotropic even though the medium is isotropic with respect to its rest frame[12, 13]. This is clear, because the isotropy of the medium is not preserved under a Lorentz transformation. Furthermore the electromagnetic features of the moving medium in direction of its motion are different from those in any other directions.

3. A medium which is only spatially inhomogeneous with respect to its rest frame is both spatially and temporarily inhomogeneous when it moves. Therefore a moving medium will be in general both spatially and temporarily dispersive even if it can be assumed to respond locally when it is at rest.

To obtain the relativistically covariant form of the constitutive equation of the medium which satisfies the above mentioned properties, we write the Euler-Lagrange equation for the tensor $Y^{\alpha\beta}(\omega, x)$

$$\square^2 Y^{\alpha\beta}(\omega, x) + \omega^2 Y^{\alpha\beta}(\omega, x) = -f^{\mu\nu\alpha\beta}(\omega, x) \partial_\mu A_\nu(x) \quad (4)$$

By solving this equation for the tensor $Y^{\alpha\beta}$ in terms the retarded Green function [11, 14] and substituting the obtained solution in the definition of the polarization tensor (3), the covariant form of the constitutive equation of the the medium relating the polarization tensor $K^{\mu\nu}$ to the the electromagnetic field strength tensor $F^{\mu\nu}$ is found as

$$K^{\mu\nu}(x) = K_N^{\mu\nu}(x) + \int_{-\infty}^{+\infty} d^4 x' \chi^{\mu\nu\alpha\beta}(x, x') F_{\alpha\beta}(x') \quad (5)$$

where $\chi^{\mu\nu\alpha\beta}(x, x')$ is the susceptibility tensor of the medium which is anti-symmetric with respect to the first two indices μ, ν and also with respect to the last two indices α, β and is given by

$$\begin{aligned} \chi^{\mu\nu\alpha\beta}(x, x') &= \frac{1}{8\pi} \int_0^\infty d\omega G(\omega, x - x') f^{\mu\nu\delta\sigma}(\omega, x) f^{\alpha\beta}_{\delta\sigma}(\omega, x') \\ &= -\frac{\vartheta(t - t')}{64\pi^4} \int_D d^4 q \sin[q \cdot (x - x')] \frac{f^{\mu\nu\delta\sigma}(\sqrt{q^2}, x) f^{\alpha\beta}_{\delta\sigma}(\sqrt{q^2}, x')}{2\sqrt{q^2}} \end{aligned} \quad (6)$$

Here ϑ is the step function , the integration domain D is the upper half cone $q_0^2 - \vec{q}^2 \geq 0$, $q_0 \geq 0$ and $G(\omega, x - x')$ is the retarded Green function for Eq. (4) satisfying

$$[\square^2 + \omega^2] G(\omega, x - x') = -\delta^{(4)}(x - x') \quad (7)$$

In the constitutive equation (5), $K_N^{\mu\nu}$ is the noise polarization tensor which is related to the solution of the homogeneous equation $[\square^2 + \omega^2]Y_N^{\alpha\beta} = 0$ as follows

$$K_N^{\mu\nu}(x) = \frac{1}{4\pi} \int_0^\infty d\omega f^{\mu\nu\delta\sigma}(\omega, x) Y_{N\delta\sigma}(\omega, x) \quad (8)$$

From the solution of the wave equation (4) which is written as

$$Y^{\mu\nu}(\omega, x) = Y_N^{\mu\nu}(\omega, x) + \int_{-\infty}^{+\infty} d^4x' G(\omega, x - x') f^{\mu\nu\alpha\beta}(\omega, x') \partial_\alpha A_\beta(x') \quad (9)$$

since the retarded Green function $G(\omega, x - x')$ vanishes in the limit $t \rightarrow -\infty$ it is clear that the noise fields $Y_N^{\alpha\beta}(\omega, x)$ and $K_N^{\mu\nu}(x)$ are the asymptotic forms of $Y^{\alpha\beta}(\omega, x)$ and $K^{\mu\nu}$ for very large negative times, respectively. The noise tensor field $Y_N^{\alpha\beta}(\omega, x)$ can be expanded in terms of the plane waves as follows

$$Y_N^{\alpha\beta}(\omega, x) = \frac{1}{\sqrt{(2\pi)^3}} \int_{-\infty}^{+\infty} d^3k [B^{\alpha\beta}(\omega, \vec{k}) e^{i\vec{k}\cdot\vec{x} - i\sqrt{k^2 + \omega^2}t} + C.C] \quad (10)$$

In classical electrodynamics the tensor fields $B^{\alpha\beta}(\omega, \vec{k})$ are numerical functions which can be obtained in terms of the asymptotic forms of $Y^{\alpha\beta}(\omega, x)$ and $\frac{\partial Y^{\alpha\beta}(\omega, x)}{\partial t}$ as

$$B^{\alpha\beta}(\omega, \vec{k}) e^{-i\sqrt{k^2 + \omega^2}t} = \frac{1}{2\sqrt{(2\pi)^3}} \int_{-\infty}^{+\infty} d^3x [Y_N^{\alpha\beta}(\omega, x) + \frac{i}{\sqrt{k^2 + \omega^2}} \frac{\partial Y_N^{\alpha\beta}(\omega, x)}{\partial t}] e^{-i\vec{k}\cdot\vec{x}} \quad (11)$$

In Quantum electrodynamics which is discussed in the next section the tensor fields $B^{\alpha\beta}(\omega, \vec{k})$ are operator valued functions that act on the Fock space of the total system.

3 Canonical quantization

Having the Lagrangian of the total system given by (1), one can compute the canonical conjugate momentum densities of the system as

$$\begin{aligned} -\Pi^\nu &= \frac{\partial L}{\partial(\partial_0 A_\nu)} = \partial^0 A^\nu - 4\pi K^{0\nu} \\ Q^{\alpha\beta}(\omega, x) &= \frac{\delta L}{\delta(\partial_0 Y_{\alpha\beta}(\omega, x))} = \partial^0 Y^{\alpha\beta}(\omega, x) \end{aligned} \quad (12)$$

According to the standard canonical quantization method, to quantize the total system the following equal-time commutation relations are imposed on the dynamical fields $A^\nu, Y^{\alpha\beta}(\omega, x)$ and their conjugate variables

$$\left[A^\mu(\vec{x}, t), -\Pi^\nu(\vec{x}', t) \right] = i\hbar g^{\mu\nu} \delta^{(3)}(\vec{x} - \vec{x}') \quad (13)$$

$$\left[Y^{\alpha\beta}(\omega, \vec{x}, t), Q^{\mu\nu}(\omega', \vec{x}', t) \right] = i\hbar [g^{\alpha\mu} g^{\beta\nu} - g^{\alpha\nu} g^{\beta\mu}] \delta(\omega - \omega') \delta^{(3)}(\vec{x} - \vec{x}') \quad (14)$$

The canonical momenta (12) and the total Lagrangian (1) can be used to write the Hamiltonian of the total system as

$$\begin{aligned} H(t) = & \int_{-\infty}^{+\infty} d^3x \left[\frac{(\Pi^\alpha - K^{0\alpha})(\Pi_\alpha - K_{0\alpha})}{2} \right] \\ & + \int_{-\infty}^{+\infty} d^3x \left[-\frac{1}{2} \partial_l A_\alpha \partial^l A^\alpha + 4\pi \partial_l A_\alpha K^{l\alpha} \right] \\ & + \int_{-\infty}^{+\infty} d^3x \int_0^\infty d\omega \left[\frac{1}{2} Q_{\alpha\beta}(\omega, \vec{x}, t) Q^{\alpha\beta}(\omega, \vec{x}, t) \right] \\ & - \frac{1}{2} \int_{-\infty}^{+\infty} d^3x \left[\partial_l Y_{\alpha\beta}(\omega, \vec{x}, t) \partial^l Y^{\alpha\beta}(\omega, \vec{x}, t) \right] \\ & + \int_{-\infty}^{+\infty} d^3x \left[\frac{1}{2} \omega^2 Y_{\alpha\beta}(\omega, \vec{x}, t) Y^{\alpha\beta}(\omega, \vec{x}, t) \right] \end{aligned} \quad (15)$$

where the sum is done over α from 0 to 3 and over l from 1 to 3. It is easy to show that, in the Heisenberg picture, the combination of the Heisenberg equations of the conjugate variables A^ν and $-\Pi^\nu$ leads to the covariant wave equation (2). Also using the Heisenberg equations of the conjugate variables $Y^{\alpha\beta}(\omega, x)$ and $Q^{\alpha\beta}(\omega, x)$ gives us Eq.(4). Now substituting $K^{\mu\nu}$ from (5), into (2) reads to the equation of motion of the vector potential as

$$\square^2 A^\nu - 8\pi \int_{-\infty}^{+\infty} d^4x' \partial_\mu \chi^{\mu\nu\alpha\beta}(x, x') \partial_\alpha A_\beta(x') = 4\pi \partial_\mu K_N^{\mu\nu}(x) \quad (16)$$

By introducing the following basis for the second rank antisymmetric tensors

$$\eta^{\alpha\beta}(\vec{k}, \lambda, \lambda') = \frac{\varepsilon^\alpha(\vec{k}, \lambda) \varepsilon^\beta(\vec{k}, \lambda') - \varepsilon^\alpha(\vec{k}, \lambda') \varepsilon^\beta(\vec{k}, \lambda)}{\sqrt{2}} \quad \lambda < \lambda', \quad (17)$$

the noise tensor field $Y_N^{\alpha\beta}(\omega, x)$ can be expanded in terms of the plane waves as follows

$$Y_N^{\alpha\beta}(\omega, x) = \sum_{\lambda, \lambda'=0}^3 \int_{-\infty}^{+\infty} d^3k \sqrt{\frac{\hbar}{4(2\pi)^3 \sqrt{k^2 + \omega^2}}} \times \\ [b_{\lambda\lambda'}(\omega, \vec{k}) e^{i\vec{k} \cdot \vec{x} - i\sqrt{k^2 + \omega^2}t} + h.c.] \eta^{\alpha\beta}(\lambda, \lambda', \vec{k}), \quad (18)$$

where $\varepsilon^\alpha(\vec{k}, \lambda)$ $\lambda = 0, 1, 2, 3$ are a basis for the four dimensional Minkowski space and for any vector $\vec{k}$ obeying the orthonormality and the completeness relations

$$\varepsilon^\sigma(\lambda, \vec{k}) \varepsilon_\sigma(\lambda', \vec{k}) = g_{\lambda\lambda'} \\ \sum_{\lambda=0}^3 g_{\lambda\lambda'} \varepsilon_\sigma(\lambda, \vec{k}) \varepsilon_{\sigma'}(\lambda, \vec{k}) = g_{\sigma\sigma'}. \quad (19)$$

The annihilation and creation operators $b_{\lambda\lambda'}(\omega, \vec{k})$ and $b_{\lambda\lambda'}^\dagger(\omega, \vec{k})$ in the expansion (18) are antisymmetric with respect to the indices λ, λ' . Since the noise fields $Y_N^{\alpha\beta}(\omega, x)$ and $\partial_0 Y_N^{\alpha\beta}(\omega, x)$ obey the same commutation relations (14), the commutation relations between $b_{\lambda\lambda'}(\omega, \vec{k})$ and $b_{\lambda\lambda'}^\dagger(\omega, \vec{k})$ are

$$[b_{\lambda_1\lambda_2}(\omega, \vec{k}), b_{\lambda'_1\lambda'_2}^\dagger(\omega', \vec{k}')] = [g_{\lambda_1\lambda'_1} g_{\lambda_2\lambda'_2} - g_{\lambda_1\lambda'_2} g_{\lambda_2\lambda'_1}] \delta(\vec{k} - \vec{k}') \delta(\omega - \omega'), \quad (20)$$

where the completeness and orthonormality relations

$$\sum_{\lambda, \lambda'=0}^3 g_{\lambda\lambda} g_{\lambda'\lambda'} \eta^{\alpha\beta}(\lambda, \lambda', \vec{k}) \eta^{\mu\nu}(\lambda, \lambda', \vec{k}) = g^{\alpha\mu} g^{\beta\nu} - g^{\alpha\nu} g^{\beta\mu}, \\ \eta^{\alpha\beta}(\lambda_1, \lambda'_1, \vec{k}) \eta_{\alpha\beta}(\lambda_2, \lambda'_2, \vec{k}) = g_{\lambda_1\lambda_2} g_{\lambda'_1\lambda'_2} - g_{\lambda_1\lambda'_2} g_{\lambda'_1\lambda_2} \quad (21)$$

has been used. The noise polarization tensor $K_N^{\alpha\beta}$ defined by (8) can now be expressed in terms of the ladder operators of the medium as

$$K_N^{\alpha\beta}(x) = \frac{1}{4\pi} \int_0^\infty d\omega f^{\alpha\beta\mu\nu}(\omega, x) \sum_{\lambda, \lambda'=0}^3 \int_{-\infty}^{+\infty} d^3k \sqrt{\frac{\hbar}{4(2\pi)^3 \sqrt{k^2 + \omega^2}}} \times \\ [b_{\lambda\lambda'}(\omega, \vec{k}) e^{i\vec{k} \cdot \vec{x} - i\sqrt{k^2 + \omega^2}t} + h.c.] \eta_{\mu\nu}(\lambda, \lambda', \vec{k}) \quad (22)$$

In this quantization scheme the Lorentz condition, $\partial_\mu A^\mu = 0$, as an operator identity is not compatible to the commutation relations (13). Instead

only those state vectors $|\psi\rangle$ in the Hilbert space of the total system are admissible for which the expectation value of the gauge condition is satisfied, that is $\langle\psi|\partial_\mu A^\mu|\psi\rangle = 0$ [13].

It is remarkable that the coupling tensor $f^{\mu\nu\alpha\beta}(\omega, x)$ satisfying the relation (6) for an assumed susceptibility tensor $\chi^{\mu\nu\alpha\beta}$ may not be unique. But the various choices of $f^{\mu\nu\alpha\beta}(\omega, x)$ satisfying (6) for a given susceptibility tensor are equivalent and do not change the commutation relations between the electromagnetic field operators. This becomes more clear if, using (20) and (22), we compute the commutation relations of the components of the noise polarization tensor for arbitrary points x, x' in space-time

$$\left[K_N^{\mu\nu}(x), K_N^{\alpha\beta}(x') \right] = \frac{i\hbar}{\pi} \left[\vartheta(x_0 - x'_0) \chi^{\mu\nu\alpha\beta}(x, x') - \vartheta(x'_0 - x_0) \chi^{\alpha\beta\mu\nu}(x', x) \right]. \quad (23)$$

These commutation relations remain unchanged under the various choices of the coupling tensor $f^{\mu\nu\alpha\beta}(\omega, x)$ satisfying (6).

For a moving homogeneous bulk material, that is a medium for which the tensor $\chi^{\mu\nu\alpha\beta}(x, x')$ is dependent only on the difference $x - x'$, it is clear from (6) that the coupling tensor $f^{\mu\nu\alpha\beta}(\omega, x)$ is independent of x and the wave equation (16) can be solved using the four dimensional Fourier transform technique to obtain the vector potential as

$$A_\nu(x) = \int_0^{+\infty} d\omega \int_{-\infty}^{+\infty} d^3k \sum_{\lambda, \lambda'=0}^3 \sqrt{\frac{\hbar}{2(2\pi)^3 \sqrt{k^2 + \omega^2}}} \times \\ \left[Z_{\nu\alpha\beta}^*(\vec{k}, \omega) b_{\lambda\lambda'}(\omega, \vec{k}) e^{i\vec{k} \cdot \vec{x} - i\sqrt{k^2 + \omega^2} t} + h.c \right] \eta^{\alpha\beta}(\lambda, \lambda', \vec{k}) \quad (24)$$

$$Z_{\nu\alpha\beta}(\vec{k}, \omega) = f^{\mu\sigma}_{\alpha\beta}(\omega) \left[i q_\mu L_{\nu\sigma}^{-1}(q) \right]_{q=(\vec{k}, \sqrt{k^2 + \omega^2})} \quad (25)$$

where

$$L_{\nu\sigma}(q) = \left[-q_\mu q^\mu \delta_{\nu\sigma} + 8\pi q^\mu q^\alpha \underline{\chi}_{\mu\nu\alpha\sigma}(q) \right] \quad (26)$$

and $\underline{\chi}^{\mu\nu\alpha\beta}(q)$ is the four dimensional Fourier transform of the tensor $\chi^{\mu\nu\alpha\beta}$. Applying a boost transformation on the susceptibility tensor $\chi^{\mu\nu\alpha\beta}$, one can show that a medium which is isotropic in its rest frame, becomes an anisotropic one when it moves with a constant velocity $\vec{v}$. In this case the electromagnetic field operators are obtained dependently on the magnitude and the direction of the velocity of the moving medium.

References

- [1] F. Kheirandish, M. Amooshahi, Phys. Rev. A 74, 042102 (2006).
- [2] M. Amooshahi, F. Kheirandish, Phys. Rev. A 76, 062103 (2007).
- [3] M. Amooshahi, F. Kheirandish, Modd. Phys. Lett. A 23, No.26,2163(2008).
- [4] M. Amooshahi, F. Kheirandish, J. Phys. A: Math. Theor. 41, No. 27, 275402(2008).
- [5] M. Amooshahi, Arxive, 0810 (2008)2695.
- [6] B. Huttner S. M. Barnett, Phys. Rev. A 46, 4306(1992).
- [7] L. G. Sutorp, M. Wubs, Phys. Rev. A 70, 013816(2004).
- [8] L. G. Sutorp, J. Phys. A: Math. Theor. 40, No. 27, No. 13, 3697(2007).
- [9] W. Greiner, J. Reinhardt, Field Quantization, Springer-Verlag Berlin Heidelberg (1996).
- [10] H. C. Chen, Theory of Electromagnetic Waves, McGraw-Hill, New York (1983).
- [11] L. D. Landau, E. M. Lifshitz, Electrodynamics of Continuous Media, Pergamon Press, Oxford, New york (1960).
- [12] R. Matloob, Phys. Rev. A 71, 062105 (2005).
- [13] R. Matloob, Phys. Rev. A 72, 062103(2005) .
- [14] J. D. Jackson, Classical Electrodynamics, John Wiley and Sons, New york (1999).